


Problem 1. Let X be a random variable with density given by

$$f(x) = \begin{cases} c(1-x^2) & -1 < x < 1 \\ 0 & \text{otherwise.} \end{cases}$$

- Find c
- Find the CDF of X .
- Find $P(-.5 \leq X < 0.5)$.
- Find $E[X]$ and $V(X)$.



$$\begin{aligned} \textcircled{a} \quad 1 &= \int_{-1}^1 c(1-x^2) dx = 2c \int_0^1 (1-x^2) dx \\ &= 2c \left(x - \frac{1}{3}x^3 \Big|_0^1 \right) \\ &= \frac{4}{3}c, \quad \text{so } c = \frac{3}{4} \end{aligned}$$

$$\begin{aligned} \textcircled{b} \quad \text{when } -1 < x < 1, \quad F(x) &= P(\bar{X} \leq x) = \int_{-1}^x c(1-t^2) dt \\ &= c \left(t - \frac{1}{3}t^3 \Big|_{-1}^x \right) \\ &= c \left[\left(x - \frac{1}{3}x^3 \right) - \left(-1 + \frac{1}{3} \right) \right] \\ &= \frac{3}{4} \left(x - \frac{1}{3}x^3 \right) + \frac{1}{2}. \end{aligned}$$

$$\text{So } F(x) = \begin{cases} 0 & x \leq -1 \\ \frac{3}{4} \left(x - \frac{1}{3}x^3 \right) + \frac{1}{2} & -1 < x < 1 \\ 1 & x \geq 1 \end{cases}$$

$$\begin{aligned} \textcircled{c} \quad P(-\frac{1}{2} \leq \bar{X} \leq \frac{1}{2}) &= F(\frac{1}{2}) - F(-\frac{1}{2}) \\ &= \frac{3}{4} \left(\frac{1}{2} - \frac{1}{24} \right) - \frac{3}{4} \left(-\frac{1}{2} + \frac{1}{24} \right) \\ &= \frac{11}{16} \end{aligned}$$

$$\begin{aligned} \textcircled{d} \quad E[\bar{X}] &= \int_{-1}^1 cx(1-x^2) dx = 0 \\ V(\bar{X}) &= E[\bar{X}^2] - E[\bar{X}]^2 = \int_{-1}^1 cx^2(1-x^2) dx \\ &= 2c \int_0^1 (x^2 - x^4) dx \\ &= 2c \left(\frac{1}{3}x^3 - \frac{1}{5}x^5 \right) \Big|_0^1 \\ &= \frac{4c}{15} = \frac{4}{15} \left(\frac{3}{4} \right) = \frac{1}{5} \end{aligned}$$

Problem 2. Suppose that X is a random variable with CDF given by

$$F(x) = \begin{cases} 1 - \frac{1}{x^2} & x > 1 \\ 0 & \text{otherwise.} \end{cases}$$

- Find the median of X .
- Find $P(X \geq 1.5)$
- Find $P(3 \leq X \leq 4)$
- Find the density of X .

(a) find m so that $F(m) = P(\bar{X} \leq m) = \frac{1}{2}$:

$$1 - \frac{1}{m^2} = \frac{1}{2} \Rightarrow \frac{1}{m^2} = \frac{1}{2} \Rightarrow m^2 = 2 \Rightarrow m = \sqrt{2}$$

(b) $P(\bar{X} \geq 1.5) = 1 - P(\bar{X} < 1.5)$
 $= 1 - F(1.5)$
 $= \frac{1}{(1.5)^2} = \frac{4}{9}$

(c) $P(3 \leq \bar{X} \leq 4) = P(\bar{X} \leq 4) - P(\bar{X} < 3)$
 $= F(4) - F(3)$
 $= \frac{1}{3^2} - \frac{1}{4^2} = \frac{7}{144}$

(d) $f(x) = F'(x) = \begin{cases} \frac{2}{x^3} & x > 1 \\ 0 & x \leq 1. \end{cases}$

Problem 3. Someone receives on average 5 text messages per hour. Let X denote the time until they receive a message.

- What distribution should be used to model X ? What parameter(s) does it have?
- Find the expected time until receiving a message.
- Find the probability that it takes between 10 and 20 minutes before receiving a message.
- Find the probability that it takes less 30 minutes to receive a message.
- Suppose no message has been received after 30 minutes. Find the probability that still no message has been received after 5 more minutes.

(a) $\bar{X} \sim \text{Exp}(5)$ (in hours) or $\bar{X} \sim \text{Exp}(1/12)$ (in minutes)

(b) $1/5$ hour or 12 minutes

(c) $P(10 \leq \bar{X} \leq 20) = F(20) - F(10)$
 $= (1 - e^{-20\lambda}) - (1 - e^{-10\lambda}) = e^{-10/12} - e^{-20/12}$

(d) $P(\bar{X} \leq 30) = F(30) = 1 - e^{-30\lambda} = 1 - e^{-30/12}$

(e) $P(\bar{X} > 35 | \bar{X} > 30) = P(\bar{X} > 5) = 1 - P(\bar{X} \leq 5) = 1 - F(5) = e^{-5/12}$

Problem 4. Suppose you are working at a blood bank where people donate blood. Type O blood is one of the best to be donated since it can be used for many people. Approximately 42% of people have type O blood.

- Find the probability that it takes exactly 15 patients until you've gotten 1 with type O blood.
- Find the probability that it takes exactly 15 patients until you've gotten 3 with type O blood.
- On average how many patients have to come in until you've gotten 3 with type O blood?

Ⓐ Let $\bar{X} \sim \text{Geom}(0.42)$. $P(\bar{X}=15) = (0.58)^{14} (0.42)$

Ⓑ Let $\bar{I} \sim \text{Neg Bin}(3, 0.42)$ $P(\bar{I}=15) = \binom{14}{2} (0.58)^{12} (0.42)^3$

Ⓒ $E[\bar{I}] = \frac{3}{0.42}$

Problem 5. For the following situations describe the distribution, including parameters, of the given random variables. Give the most reasonable distribution for the situation.

- Every day there is a 10% chance that Rick will receive no mail. Let X be the number of times he receives mail over the next 5 days.
- Shawna is playing craps at the casino. The probability of winning craps is about 0.49. She will keep playing until she wins. Let X be the number of times she will play.
- Of 100 raffle tickets available, 30 are marked for a prize by the presence of a blue dot. Courtney picks up 10 raffle tickets at random for her co-workers. Let X be the number of raffle tickets Courtney has that are marked for a prize.
- Rainer is listening to a music station which he isn't used to, and finds that he recognizes about 30% of the songs that they play. He plans to keep listening to songs until he hears 4 songs that he knows. Let X be the number of total songs that Rainer has to listen to in order to hear 4 songs that he knows.

Ⓐ $\text{Bin}(5, 0.9)$

Ⓑ $\text{Geom}(0.49)$

Ⓒ $\text{Hyper Geom}(30, 100, 10)$

\swarrow distinguished portion of population r
 \uparrow population size N
 \nwarrow sample size n

Ⓓ $\text{Neg Bin}(4, 0.3)$

Problem 6. Faith is handing out treats to her cat, Mysteria. She has been randomly generating X , the number of treats using a binomial distribution with $n = 3$ and $p = 0.4$. Malcolm says that it's unfair that Mysteria might get 0 treats, so he suggests 0 be ruled out as an option, and those values all turned into 1's, so that the cat always gets at least one treat. Denote Malcolm's new random variable for the number of treats by U .

- Find the moment generating function of U .
- Use the moment generating function of U to compute $E[U]$.

$$\textcircled{a} \quad P(U=k) = \begin{cases} (0.6)^3 + 3(0.4)(0.6)^2 & k=1 \\ 3(0.4)^2(0.6) & k=2 \\ (0.4)^3 & k=3 \end{cases} = \begin{cases} 81/125 & k=1 \\ 36/125 & k=2 \\ 8/125 & k=3 \end{cases}$$

$$m(t) = E[e^{tU}] = \frac{81}{125} e^t + \frac{36}{125} e^{2t} + \frac{8}{125} e^{3t}$$

$$\textcircled{b} \quad m'(t) = \frac{81}{125} e^t + \frac{72}{125} e^{2t} + \frac{24}{125} e^{3t}$$

$$E[U] = m'(0) = \frac{81}{125} + \frac{72}{125} + \frac{24}{125} = \frac{177}{125}$$

Problem 7. Let X be the value of the first die and Y the sum of the values when two dice are rolled. Compute the moment generating functions of X and Y .

$$\bar{X} \sim \text{Unif}(\{1, 2, 3, 4, 5, 6\})$$

$$m_{\bar{X}}(t) = E[e^{t\bar{X}}] = \frac{1}{6}(e^t + e^{2t} + e^{3t} + e^{4t} + e^{5t} + e^{6t})$$

$$\bar{Y} = \bar{X}_1 + \bar{X}_2 \quad \text{where } \bar{X}_1, \bar{X}_2 \sim \text{Unif}(\{1, 2, 3, 4, 5, 6\}) \text{ are i.i.d.}$$

$$\begin{aligned} m_{\bar{Y}}(t) &= E[e^{t\bar{Y}}] = E[e^{t(\bar{X}_1 + \bar{X}_2)}] \\ &= E[e^{t\bar{X}_1}] E[e^{t\bar{X}_2}] \\ &= m_{\bar{X}}(t)^2 \\ &= \frac{1}{36} (e^t + e^{2t} + e^{3t} + e^{4t} + e^{5t} + e^{6t})^2 \end{aligned}$$

Problem 8. A bag contains 3 red, 5 green, and 7 blue balls. A sample of 2 balls is drawn with replacement. Let X be the number of red balls in the sample and let Y be the number of green balls in the sample.

- Find the joint probability mass function of X and Y .
- Find $P(X > Y)$.
- Find the marginal distributions of X and Y .
- Find $E[X + Y]$.

Ⓐ

	$\bar{Y}=0$	$\bar{Y}=1$	$\bar{Y}=2$	
$\bar{X}=0$	$(\frac{7}{15})^2$	$2(\frac{5}{15})(\frac{7}{15})$	$(\frac{5}{15})^2$	$\frac{144}{225}$
$\bar{X}=1$	$2(\frac{3}{15})(\frac{7}{15})$	$2(\frac{3}{15})(\frac{5}{15})$	0	$\frac{72}{225}$
$\bar{X}=2$	$(\frac{3}{15})^2$	0	0	$\frac{9}{225}$
	$\frac{100}{225}$	$\frac{100}{225}$	$\frac{25}{225}$	

Ⓑ $P(\bar{X} > \bar{Y}) = P(\bar{X}=1, \bar{Y}=0) + P(\bar{X}=2, \bar{Y}=0)$

$$= \frac{42}{225} + \frac{9}{225} = \frac{51}{225}$$

Ⓒ

$$P(\bar{X}=k) = \begin{cases} \frac{144}{225} & k=0 \\ \frac{72}{225} & k=1 \\ \frac{9}{225} & k=2 \end{cases} \quad P(\bar{Y}=k) = \begin{cases} \frac{100}{225} & k=0 \\ \frac{100}{225} & k=1 \\ \frac{25}{225} & k=2 \end{cases}$$

Ⓓ $E[\bar{X} + \bar{Y}] = E[\bar{X}] + E[\bar{Y}]$

$$= \left(\frac{72}{225} + \frac{18}{225} \right) + \left(\frac{100}{225} + \frac{50}{225} \right)$$

$$= \frac{240}{225}$$

Problem 9. Let

$$E[X] = 5, E[X^2] = 27.5, E[X^3] = 162.5, E[X^4] = 1017.5,$$

$$E[Y] = 6, E[Y^2] = 39.6, E[Y^3] = 281.52, E[Y^4] = 2128.176.$$

Suppose X and Y are independent. Find the following quantities.

- $V(X - Y^2)$
- $E[X^2 Y^3]$
- $V(X^2 Y^2)$

$$\begin{aligned} \textcircled{a} \quad V(X - Y^2) &= V(X) + V(Y^2) \\ &= (27.5 - 5^2) + (2128.176 - 39.6^2) \end{aligned}$$

$$\begin{aligned} \textcircled{b} \quad E[X^2 Y^3] &= E[X^2] E[Y^3] \\ &= (27.5)(281.52) \end{aligned}$$

$$\begin{aligned} \textcircled{c} \quad V(X^2 Y^2) &= E[(X^2 Y^2)^2] - E[X^2 Y^2]^2 \\ &= E[X^4] E[Y^4] - E[X^2]^2 E[Y^2]^2 \\ &= (1017.5)(2128.176) - (27.5)^2 (39.6)^2 \end{aligned}$$