

§ 3.5 Poisson distribution

When modeling with the binomial distribution,
we require a fixed number of trials. However,
consider counting:

- number of babies born in a hospital in 1 day
- number of emails received in 1 hour
- number of highway accidents in 1 year

These scenarios involve counting during a fixed time period
but the notion of independent trials doesn't make sense.

Instead, we model them using the Poisson distribution.

Def A random variable $\bar{X}$ has the Poisson distribution
with parameter $\lambda > 0$ if its range is $\{0, 1, 2, \dots\}$ and

$$P(\bar{X} = k) = e^{-\lambda} \frac{\lambda^k}{k!} \quad \text{for } k = 0, 1, 2, \dots$$

Shorthand: $\bar{X} \sim \text{Pois}(\lambda)$.

Remark: the parameter λ is the average count (or
we often use the term "number of arrivals" instead of "count")
over the fixed time period

Note $\sum_{k=0}^{\infty} e^{-\lambda} \frac{\lambda^k}{k!} = e^{-\lambda} \sum_{k=0}^{\infty} \frac{\lambda^k}{k!} = e^{-\lambda} \cdot e^{\lambda} = 1$

since $e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!}$ for all $x \in \mathbb{R}$ (Taylor series)

Example Suppose someone receives on average 10 emails per hour. We model the number of emails in 1 hour as a random variable $\bar{X}$ with the Poisson distribution.

- ① Find the probability they receive exactly 9 emails in 1 hour.

$$\bar{X} \sim \text{Pois}(10) \quad \text{and} \quad P(\bar{X}=9) = e^{-10} \frac{10^9}{9!}$$

- ② Find the probability they receive exactly 21 calls

in 2 hours. Let $\bar{Y} \sim \text{Pois}(20)$, $P(\bar{Y}=21) = e^{-20} \frac{20^{21}}{21!}$

Example Suppose someone receives 2 calls per hour on average. What is the probability that over 24 hours they receive exactly 50 calls? at least 50 calls?

Let $\bar{X} \sim \text{Pois}(48)$ count the number of calls received over 24 hours. Then

$$P(\bar{X}=50) = e^{-48} \frac{48^{50}}{50!}$$

$$P(\bar{X} \geq 50) = 1 - P(\bar{X} \leq 49)$$

compute this with R

R commands if $\bar{X} \sim \text{Pois}(\lambda)$, then

- $P(\bar{X}=k)$ can be computed in R with

$$\text{dpois}(k, \lambda)$$

- $P(\bar{X} \leq k)$ with $\text{ppois}(k, \lambda)$


- we can get an i.i.d. sample of size k with

$$\text{rpois}(k, \lambda).$$

Derivation of the Poisson distribution

Suppose on average λ babies are born over 1 day.

Let's break up the day into n pieces of length $\frac{1}{n}$.



- On average, in any of these subintervals $\frac{\lambda}{n}$ babies are born.
- Moreover, if $\bar{X}_1, \dots, \bar{X}_n$ denotes the number of babies born in the first, second, etc. subinterval, then when n is big enough each of these $\bar{X}_k$ could only take values in $\{0, 1\}$, (ie. it's very unlikely to have 2 or more babies born simultaneously)
- So $\bar{X}_1, \dots, \bar{X}_n$ are i.i.d. $\text{Ber}(p)$ with $p = \frac{\lambda}{n}$.

- Since $\bar{X} = \lim_{n \rightarrow \infty} \bar{X}_1 + \dots + \bar{X}_n$ and $\bar{X}_1 + \dots + \bar{X}_n \sim \text{Bin}(n, \frac{\lambda}{n})$

$$\begin{aligned} P(\bar{X} = k) &= \lim_{n \rightarrow \infty} \binom{n}{k} \left(\frac{\lambda}{n}\right)^k \left(1 - \frac{\lambda}{n}\right)^{n-k} \\ &= e^{-\lambda} \frac{\lambda^k}{k!} \quad (\text{using calculus}) \end{aligned}$$

$$\lim_{n \rightarrow \infty} \binom{n}{k} \left(\frac{\lambda}{n}\right)^k \left(1 - \frac{\lambda}{n}\right)^{n-k} = \lim_{n \rightarrow \infty} \frac{n!}{k!(n-k)!} \frac{\lambda^k}{n^k} \left(1 - \frac{\lambda}{n}\right)^{n-k}$$

$$= \frac{\lambda^k}{k!} \lim_{n \rightarrow \infty} \frac{n!}{(n-k)!} \frac{1}{n^k} \left(1 - \frac{\lambda}{n}\right)^n \underbrace{\left(1 - \frac{\lambda}{n}\right)^{-k}}_{\rightarrow 1}$$

$$= \frac{\lambda^k}{k!} \lim_{n \rightarrow \infty} \frac{n(n-1)(n-2)\dots(n-k+1)}{\underbrace{n^k}_{\rightarrow 1}} \left(1 - \frac{\lambda}{n}\right)^n$$

$$= \frac{\lambda^k}{k!} \lim_{n \rightarrow \infty} \left(1 - \frac{\lambda}{n}\right)^n$$

$$= e^{-\lambda} \frac{\lambda^k}{k!}$$

Problem 1. Customers arrive at a checkout counter in a department store according to a Poisson distribution at an average of 7 per hour.

- Find the probability
 - exactly five customers arrive in the next hour
 - no more than three customers arrive in the next hour
 - at least two customers arrive in the next hour
- Answer the same questions but change "next hour" to "next 30 minutes."

$$\textcircled{a} \quad \bar{X} \sim \text{Pois}(7)$$

$$P(\bar{X} = 5) = e^{-7} \frac{7^5}{5!} \approx 0.1277167$$

$$P(\bar{X} \leq 3) \approx 0.08176542$$

$$P(\bar{X} \geq 2) \approx 0.9927049$$

$$\textcircled{b} \quad \bar{X} \sim \text{Pois}(3.5), \text{ see right for values}$$

Part a:

```

```{r}
dpois(5, 7)
ppois(3, 7)
1-ppois(1, 7)
```

[1] 0.1277167
[1] 0.08176542
[1] 0.9927049

```

Part b:

```

```{r}
dpois(5, 3.5)
ppois(3, 3.5)
1-ppois(1, 3.5)
```

[1] 0.1321686
[1] 0.5366327
[1] 0.8641118

```

Problem 2. A telemarketing company has found that the probability of making a sale when calling someone on the telephone is approximately 0.0002. If the salesperson contacts 10000 prospects, what is the probability of making between 2 and 5 sales?

- Calculate this using the binomial distribution. What are n and p ?
- Let $Y \sim \text{Pois}(np)$. Compute $P(2 \leq Y \leq 5)$ and compare with your previous answer.

$$\textcircled{a} \quad \bar{X} \sim \text{Bin}(10000, 0.0002)$$

$$P(2 \leq \bar{X} \leq 5) \approx 0.5774684$$

$$\textcircled{b} \quad \bar{Y} \sim \text{Pois}(2)$$

$$P(2 \leq \bar{Y} \leq 5) \approx 0.5774305$$

Problem 2

Part a:

```

```{r}
pbinom(5, 10000, 0.0002) - pbinom(1, 10000, 0.0002)
```

[1] 0.5774684

```

Part b:

```

```{r}
ppois(5, 10000*0.0002) - ppois(1, 10000*0.0002)
```

[1] 0.5774305

```