

§1.8 Problem Solving Strategies (complements, inclusion-exclusion)

Example We roll 4 dice. Find the probability of getting at least one 6.

Let A be the event of rolling at least one 6.

Then A^c is the event of rolling no 6.

Let Ω be the sample space.

$$\text{Then } P(A) = 1 - P(A^c) = 1 - \frac{|A^c|}{|\Omega|} = 1 - \frac{5^4}{6^4}$$

General Rule When computing probabilities of "at least one" events, consider whether the probability of the complementary event is easier.

Theorem (inclusion-exclusion) Let $A, B, C \subseteq \Omega$ be events.

$$\begin{aligned} \text{Then } P(A \cup B \cup C) &= P(A) + P(B) + P(C) \\ &\quad - P(AB) - P(AC) - P(BC) \\ &\quad + P(ABC) \end{aligned}$$

Example 18 students live together in a dorm

7 of them are in a math class

10 in bio

10 in CS

3 of them in a math class and a bio class

4 in math and CS

5 in bio and CS

1 in a math class and a bio class and a CS class

Find the probability a randomly chosen student is
in neither a math nor bio nor CS class.

Let M be the event a student is in a math class

B the event they're in a bio class,

C the event they're in a CS class.

Then $P(M) = \frac{7}{18}$, $P(B) = \frac{10}{18}$, $P(C) = \frac{10}{18}$

$P(MB) = \frac{3}{18}$, $P(MC) = \frac{4}{18}$, $P(BC) = \frac{5}{18}$,

$P(MBC) = \frac{1}{18}$.

$$P((M \cup B \cup C)^c) = 1 - P(M \cup B \cup C)$$

$$= 1 - [P(M) + P(B) + P(C) \\ - P(MB) - P(MC) - P(BC) \\ + P(MBC)]$$

$$= 1 - \frac{16}{18} = \frac{1}{9}.$$

Problem 1. Consider the random experiment of rolling a die 5 times. Find the probability of each of the following events.

- All 5 rolls land on the same value.
- All 5 rolls land on different values.
- At least 1 of the rolls lands on 4.
- At least 2 of the rolls land on 4.

$$\textcircled{a} \frac{6}{6^5} \quad \textcircled{b} \frac{6!}{6^5}$$

$$\textcircled{c} 1 - \frac{5^5}{6^5} \quad \textcircled{d} 1 - \frac{5^5}{6^5} - \frac{\binom{5}{1} 5^4}{6^5}$$

Problem 2. A certain town has 3 newspapers: A , B , and C . The proportions of townspeople who read these papers are as follows:

- A : 10 percent, B : 30 percent, C : 5 percent
- A and B : 8 percent, A and C : 2 percent, B and C : 4 percent
- all 3: 1 percent

A person is chosen at random, everyone equally likely. Find the probability that they read

- at least one newspaper.
- no newspaper.
- at least two newspapers.
- exactly one newspaper.

Let A, B, C be the events that a person likes newspapers A, B, C respectively.

$$\begin{aligned} \textcircled{a} P(A \cup B \cup C) &= P(A) + P(B) + P(C) - P(AB) - P(AC) - P(BC) + P(ABC) \\ &= 0.10 + 0.30 + 0.05 - 0.08 - 0.02 - 0.04 + 0.01 \\ &= 0.32 \end{aligned}$$

$$\textcircled{b} P((A \cup B \cup C)^c) = 1 - 0.32 = 0.68$$

$$\begin{aligned} \textcircled{c} P(AB \cup AC \cup BC) &= P(AB) + P(AC) + P(BC) \\ &\quad - P(ABC) - P(ABC) - P(ABC) + P(ABC) \\ &= 0.08 + 0.02 + 0.04 - 0.02 \\ &= 0.12 \end{aligned}$$

$$\textcircled{d} P((A \cup B \cup C)(AB \cup AC \cup BC)^c) = 0.32 - 0.12 = 0.20$$

Problem 3. Consider the experiment of rolling a die 5 times with sample space Ω . Let C be the event that at least one of the 5 rolls lands on 4. For each $k = 1, \dots, 5$, let

- A_k be the event that roll k lands on 4.
- B_k be the event that among the 5 rolls, exactly k of them land on 4.

Let B_0 be the event no rolls land on 4. State whether each of the following is true or false.

- $B_0 = C^c$ **T**
- $\Omega = B_0 \cup B_1 \cup B_2 \cup B_3 \cup B_4 \cup B_5$ **T**
- $C = A_1 \cup A_2 \cup A_3 \cup A_4 \cup A_5$ **T**
- $A_1 \cup A_2 \cup A_3 \cup A_4 \cup A_5 = B_1 \cup B_2 \cup B_3 \cup B_4 \cup B_5$ **T**
- $P(C) = P(A_1) + P(A_2) + P(A_3) + P(A_4) + P(A_5)$ **F**
- $P(C) = P(B_1) + P(B_2) + P(B_3) + P(B_4) + P(B_5)$ **T**